arXiv:1404.6481v2 [math.CV] 8 May 2014

THE KOBAYASHI BALLS OF (ℂ-)CONVEX DOMAINS

NIKOLAI NIKOLOV AND MARIA TRYBUŁA

Abstract. A pure geometric description of the Kobayashi balls of (ℂ-)convex domains is given in terms of the so-called minimal basis.

1. Introduction and results

Let D be a domain in $\mathbb{C}^n$. Denote by c_D and l_D the Carathéodory distance and the Lempert function of D, respectively:

$$c_D(z,w) = \sup\{\tanh^{-1}|f(w)| : f \in \mathcal{O}(D,\mathbb{D}),\ \text{ with } f(z)=0\},$$

$$l_D(z,w) = \inf\{\tanh^{-1}|\alpha| : \exists \varphi \in \mathcal{O}(\mathbb{D},D) \text{ with } \varphi(0)=z, \varphi(\alpha)=w\},$$

where $\mathbb{D}$ is the unit disc. The Kobayashi distance k_D is the largest pseudodistance not exceeding l_D.

We are interested in a description of the Kobayashi balls near boundary points of convex and, more generally, ℂ-convex domains in terms of parameters that reflect the geometry of the boundary. The first results in this direction can be found in [2, Theorems 1 and 5.1], where the strongly pseudoconvex case in $\mathbb{C}^n$ and the weakly pseudoconvex finite type case in $\mathbb{C}^2$ are discussed with applications[1] to invariant forms of Fatou type theorems (for the boundary values). The weakly pseudoconvex finite type case in $\mathbb{C}^2$, as well as the convex finite type case in $\mathbb{C}^n$, are treated in [6, Propositions 8.8 and 8.9] as byproducts of long considerations. The strongly pseudoconvex case in $\mathbb{C}^n$ and the weakly pseudoconvex finite type in $\mathbb{C}^2$ are particular cases of the pseudoconvex Levi corank one case which is considered in [3, Theorem 1.3]. The behavior of the Kobayashi balls in all the mentioned results is given in

2010 *Mathematics Subject Classification.* 32F17, 32F45.

Key words and phrases. Carathéodory distance, Kobayashi distance, convex domain, ℂ-convex domain.

The second named author is supported by international PhD programme "Geometry and Topology in Physical Models" of the Foundation for Polish Science. The paper was prepared during her visit to the Institute of Mathematics and Informatics, Bulgarian Academy of Science, October 2013 - April 2014.

[1]See also [14] for complex ellipsoids.

terms of the Levi geometry of the boundary which is assumed smooth and bounded.

Our aim is to describe the Kobayashi balls of (ℂ-)convex domains (not necessarily smooth and bounded) in terms of the so-called minimal basis (cf. [4, 9, 12]. The constants that appear depend only on the radius of the balls and the dimension of the domains. The respective proof is short and pure geometric. The obtained result covers [6, Propositions 8.8 and 8.9].

Assume that D contains no complex lines. Let $q \in D$ and $d_D(q) = \operatorname{dist}(q, \partial D)$. Choose $q^1 \in \partial D$ so that $\tau_1(q) := \|q^1 - q\| = d_D(q)$. Put $H_1 = q + \operatorname{span}(q^1 - q)^\perp$ and $D_1 = D \cap H_1$. Let $q^2 \in \partial D_1$ so that $\tau_2(q) := \|q^2 - q\| = d_{D_1}(q)$. Put $H_2 = q + \operatorname{span}(q^1 - q, q^2 - q)^\perp$, $D_2 = D \cap H_2$ and so on. Thus we get an orthonormal basis of the vectors $e_j = \dfrac{q^j - q}{\|q^j - q\|}$, $1 \le j \le n$, which is called minimal for D at q, and positive numbers $\tau_1(q) \le \tau_2(q) \le \cdots \le \tau_n(q)$ (the basis and the numbers are not uniquely determined). After rotation we may assume that $e_1, e_2, \ldots, e_n$ is the standard basis of $\mathbb{C}^n$.

Recall now that a open set D in $\mathbb{C}^n$ is said to be (cf. [1]):

• ℂ-convex if any non-empty intersection with a complex line is a simply connected domain.

• linearly (weakly linearly convex) convex if for any $a \in \mathbb{C}^n \setminus D$ ($p \in \partial D$) there exists a complex hyperplane through a which does not intersect D.

Note that convexity ⇒ ℂ-convexity ⇒ linear convexity ⇒ weak linear convexity (cf. [1, Theorem 2.3.9 ii)] for the second implication). Moreover, in the case of C^1-smooth bounded domains the last three notions coincide (cf. [1, Corollary 2.5.6].

In view of this remark and the inequalities $c_D \le k_D \le l_D$, we have the following quantitative information about the Carathéodory/Kobayashi/Lempert balls of (ℂ-)convex domains.[2]

Theorem 1. *Let D be a domain in $\mathbb{C}^n$, containing no complex lines, and $q \in D$. Assume that the standard basis of $\mathbb{C}^n$ is minimal for D at q. Let $r > 0$.*

(i) If D is weakly linearly convex, then

$$\max_{1\le j\le n} \frac{|z_j - q_j|}{\tau_j(q)} < \frac{e^{2r}-1}{n(e^{2r}+1)} \Rightarrow \sum_{j=1}^{n} \frac{|z_j - q_j|}{\tau_j(q)} < \frac{e^{2r}-1}{e^{2r}+1}$$
$$\Rightarrow z \in D \text{ and } l_D(q, z) < r.$$

[2]By the Lempert theorem, $c_D = k_D = l_D$ in the convex case, as well as in the bounded C^2-smooth ℂ-convex case (cf. [11]).

(ii) If D is convex, then $c_D(q,z) < r$ implies $\max\limits_{1\le j\le n} \dfrac{|z_j - q_j|}{\tau_j(q)} < e^{2r} - 1$.

(iii) If D is $\mathbb{C}$-convex, then $c_D(q,z) < r$ implies $\max\limits_{1\le j\le n} \dfrac{|z_j - q_j|}{\tau_j(q)} < e^{4r} - 1$.

So there exist constants $c' = c'(r,n)$ and $c'' = c''(r)$ such that

$$\mathbb{D}(q_1, c'\tau_1(q)) \times \cdots \times \mathbb{D}(q_n, c'\tau_n(q)) \subset \mathrm{kob}_D(q,r)$$
$$\subset \mathbb{D}(q_1, c''\tau_1(q)) \times \cdots \times \mathbb{D}(q_n, c''\tau_n(q)),$$

where $\mathrm{kob}_D(q,r)$ is the Kobayashi ball $\{z \in D : k(q,z) < r\}$ and $\mathbb{D}(p,r) = \{z \in \mathbb{C} : |z-p| < r\}$. By [4, Lemma 3.10], the sizes of these polydiscs are comparable (in terms of small/big constant depending on D) with the sizes of polydiscs in [3, 6] arising from the Levi geometry of the boundary. Thus Theorem 1 extends [6, Propositions 8.9].

Note also that if D is a proper $\mathbb{C}$-convex domain in $\mathbb{C}^n$ containing complex line, then it is biholomorphic to $D' \times \mathbb{C}^{n-k}$, where D' is a bounded domain in $\mathbb{C}^k$, $0 < k < n$. (cf. Proposition 3 and the preceding remark in [10]). So $\tau_k(q) < \infty = \tau_{k+1}(q)$ and it is easy to see that Theorem 1 remains true.

To prove Theorem 1, we need the planar cases of the following

Proposition 2. *(i) Let D be proper convex domain in $\mathbb{C}^n$. Then (cf.* [13, (2)])

$$c_D(z,w) \ge \frac{1}{2}\log\frac{d_D(z)}{d_D(w)}.$$

Moreover, if $n = 1$, then

$$c_D(z,w) \ge \frac{1}{2}\log\left(1 + \frac{|z-w|}{d_D(w)}\right).$$

(ii) Let D be proper $\mathbb{C}$-convex domain in $\mathbb{C}^n$. Then

$$c_D(z,w) \ge \frac{1}{4}\log\frac{d_D(z)}{d_D(w)}.$$

Moreover, if $n = 1$, then

$$c_D(z,w) \ge \frac{1}{4}\log\left(1 + \frac{|z-w|}{d_D(w)}\right).$$

The constants $1/2$ and $1/4$ are sharp as the examples $D = \mathbb{D}$ and $D = \mathbb{C}_* \setminus \mathbb{R}^+$ show. Note that in the $\mathbb{C}$-convex case the weaker estimate

$$c_D(z,w) \ge \frac{1}{4}\log\frac{d_D(z)}{4d_D(w)}$$

is contained in [13, Proposition 2]

Theorem 1 has a local version.

Proposition 3. *Let D be a domain in $\mathbb{C}^n$ whose boundary contains no affine discs through $a \in \partial D$. Assume that the standard basis of $\mathbb{C}^n$ is minimal for D at $q \in D$. Let $r > r' > 0$.*

(i) If D is weakly linearly convex near a, then

$$\max_{1\le j\le n} \frac{|z_j - q_j|}{\tau_j(q)} < \frac{e^{2r}-1}{n(e^{2r}+1)} \Rightarrow \sum_{j=1}^n \frac{|z_j - q_j|}{\tau_j(q)} < \frac{e^{2r}-1}{e^{2r}+1}$$
$$\Rightarrow z \in D \text{ and } l_D(q,z) < r.$$

for q sufficiently close to a.

(ii) If D is convex near a, then $k_D(q,z) < r'$ implies $\max\limits_{1\le j\le n} \dfrac{|z_j - q_j|}{\tau_j(q)} < e^{2r} - 1$ for q sufficiently close to a.

(iii) If D is $\mathbb{C}$-convex near a and bounded, then $k_D(q,z) < r'$ implies $\max\limits_{1\le j\le n} \dfrac{|z_j - q_j|}{\tau_j(q)} < e^{4r} - 1$ for q sufficiently close to a.

By any of the above three notions of convexity near a we mean that there exists a neighborhood U of a such that $D \cap U$ is an open set with the respective global convexity.

Note that in the convex case, as well as in the C^1-smooth $\mathbb{C}$-convex case, if ∂D contains no affine discs through a, then ∂D contains no analytic discs through a (cf. [12, Propoisition 7]).

2. Proofs

Proof of Theorem 1. (i) Since D contains the discs $\mathbb{D}(q_1, \tau_1(q)), \dots, \mathbb{D}(q_n, \tau_n(q))$ (lying in the respective coordinate complex planes), it contains their convex hull

$$C = \{\zeta \in \mathbb{C}^n : h(\zeta) = \sum_{j=1}^n \frac{|\zeta_j - q_j|}{\tau_j(q)} < 1\}$$

(cf. [12, Lemma 15]). Then

$$l_D(q,z) \le l_C(q,z) = \tanh^{-1} h(z)$$

(cf. [5, Proposition 3.1.10]) which implies (i).

Before proving (ii) and (iii) note that by ($\mathbb{C}$-)convexity and the construction of the minimal basis there exists a complex hyperplane $q^{j+1} + W_j$ through q^{j+1} that is disjoint from D, $j = 0, \dots, n-1$. It is not difficult to see that W_j is given by the equation

$$\alpha_{j,1}\zeta_1 + \dots + \alpha_{j,j}\zeta_j + \zeta_{j+1} = 0.$$

Let $\Lambda : \mathbb{C}^n \to \mathbb{C}^n$ be the linear mapping with matrix whose rows are given by the vectors $(\alpha_{j,1}, \dots, \alpha_{j,j}, 1, 0, \dots, 0)$. Set $\Lambda_q(\zeta) = q + \Lambda(\zeta - q)$. Note that $G = \Lambda_q(D)$ is a ($\mathbb{C}$-)convex domain. Denote by G_j the projection of G onto j-th coordinate plane. Then $G \subset G' = G_1 \times \cdots \times G_n$ and the product formula for the Carathéodory distance (cf. [5, Theorem 9.5]) implies that

$$c_D(q, z) \geq c_{G'}(q, \Lambda_q(z)) = \max_{1 \leq j \leq n} c_{G_j}(q_j, z_j). \tag{1}$$

Observe also that $d_{G_j}(q_j) = \tau_j(q)$.

(ii) If D is a convex domain, then G_j is a convex domain. Hence, by Proposition 2 (i),

$$c_{G_j}(q_j, z_j) \geq \frac{1}{2} \log \left(1 + \frac{|z_j - q_j|}{\tau_j(q)} \right)$$

and (ii) follows from here and (1).

(iii) If D is a $\mathbb{C}$-convex domain, then G_j is a simple connected domain (cf. [1, Theorem 2.3.6]). Hence, by Proposition 2 (ii),

$$c_{G_j}(q_j, z_j) \geq \frac{1}{4} \log \left(1 + \frac{|z_j - q_j|}{\tau_j(q)} \right)$$

and (iii) follows from here and (1).

Proof of Proposition 2. After translation and rotation, we may assume that $0 \in \partial D$ and $w = (d_D(w), 0, \dots, 0)$.

(i) We have that $D \subset \Pi^+ = \{\zeta \in \mathbb{C}^n : \operatorname{Re} \zeta_1 > 0\}$ and hence

$$c_D(z, w) \geq c_{\Pi^+}(z, w) = \tanh^{-1} \left| \frac{z_1 - w_1}{z_1 + \overline{w}_1} \right|$$

$$\geq \tanh^{-1} \frac{|z_1 - w_1|}{|z_1 - w_1| + 2 d_D(w)} = \frac{1}{2} \log \left(1 + \frac{|z_1 - w_1|}{d_D(w)} \right).$$

(ii) It follows by weak linear convexity that $D \cap \{\zeta_1 \in \mathbb{C}^n : \zeta_1 = 0\} = \varnothing$. Denote by D_1 the projection of D onto the ζ_1-plane. Let γ_G the Carathéodory metric of a domain G in $\mathbb{C}^k$:

$$\gamma_G(\zeta; X) = \sup\{|f'(\zeta) X| : f \in \mathcal{O}(G, \mathbb{D})\}, \quad \zeta \in G,\ X \in \mathbb{C}^k.$$

The Köbe 1/4 theorem implies that

$$\gamma_{D_1}(\zeta_1; e_1) \geq \frac{1}{4 d_{D_1}(\zeta_1)} \geq \frac{1}{4 |\zeta_1|}.$$

Since D_1 is a simply connected domain (cf. [1, Theorem 2.3.6]), then

$$c_D(z, w) \geq c_{D_1}(z_1, w_1) = \inf_s \int_0^1 \gamma_{D_1}(s(t); s'(t) dt \geq \frac{1}{4} \inf_s \int_0^1 \left| \frac{s'(t)}{s(t)} \right| dt,$$

where the infimum is taken over all smooth curves $s : [0,1] \to D_1$ with $s(0) = z_1$ and $s(1) = w_1$ (cf. [5]).

Set now

$$d(\zeta_1, \eta_1) = \log \max(1 + |1 - \zeta_1/\eta_1|, 1 + |1 - \eta_1/\zeta_1|).$$

It is easy to check that d is a distance on ${\mathbb{C}_*}^3$ with "derivative"

$$\lim_{\lambda \to 0} \frac{d(\zeta_1, \zeta_1 + \lambda)}{|\lambda|} = \frac{1}{|\zeta_1|}.$$

Then (cf. [5, Lemma 4.3.3) (d)])

$$\inf_s \int_0^1 \left| \frac{s'(t)}{s(t)} \right| dt \geq d(z_1, w_1)$$

and hence

$$c_D(z, w) \geq \frac{1}{4} d(z_1, w_1) \geq \frac{1}{4} \log \left(1 + \frac{|z_1 - w_1|}{d_D(w)} \right).$$

Proof of Proposition 3. (i) Using Theorem 1 (i), it is enough to show that

$$\lim_{q \to a} \tau_n(q) = 0. \tag{2}$$

Assume the contrary. Then there exists a sequence of points $(q^j) \to a$ such that $(\tau_n(q^j)) \to \varepsilon > 0$ and $(e^j) \to e$, where e^j is the last vector of the minimal basis for D at q^j. We may find a bounded neighborhood U of a such that $D \cap U$ is a weakly linearly convex open set. Shrinking ε (if necessary), it follows that the e-directional disc Δ with center q and radius ε is a limit of affine discs in $D \cap U$. Since $D \cap U$ is a taut open set (cf. [11, Proposition 1.5]), then $\Delta \subset \partial D$, a contradiction.

(ii) Having in mind Theorem 1 (ii), it is enough to show the following.

Claim 1. Let U be a neighborhood of a such that $D \cap U$ is convex. There exist neighborhoods $W \subset V \subset U$ of a such that if $q \in D \cap W$ and $k_D(q, z) < r'$, then $z \in V$ and $p k_{D \cap U}(q, z) \leq k_D(q, z)$, where $p = r'/r$.

To prove this claim, recall that k_D is the integrated form of the Kobayashi metric

$$\kappa_D(\zeta; X) = \inf\{|\alpha| : \exists \varphi \in \mathcal{O}(\mathbb{D}, D) \text{ with } \varphi(0) = z, \alpha \varphi'(0) = X\}$$

[3]Let $a, b, c \in \mathbb{C}_*$ and $d_1 = 1 - a/b$, $d_2 = 1 - b/c$, $d_3 = 1 - a/c$. We may assume that $d(a, c) = \log(1 + |d_3|)$. Then

$$d(a, b) + d(b, c) \geq \log(1 + |d_1|) + \log(1 + |d_2|)$$
$$= \log(1 + |d_1| + |d_2| + |d_3 - d_2 - d_1|) \geq \log(1 + |d_3|) = d(a, c).$$

(cf. [5]). Fix an $\varepsilon > 0$. Then we may find a smooth curve $s : [0,1] \to D$ such that $s(0) = q$, $s(1) = z$ and

$$k_D(q,z) + \varepsilon > \int_0^1 \kappa_D(s(t); s'(t))dt.$$

Since $D \cap U$ is convex and its boundary contains no affine discs through a, then a is a peak point for $D \cap U$ (cf. [8, Theorem 6]). Hence the strong localization property for the Kobayashi metric holds (cf. [7, Theorem 1 and Corollary 2]). So there exists a neighborhood $V \subset U$ of a such that

$$\kappa_D(\zeta; X) \geq p\kappa_{D\cap U}(\zeta; X), \quad \zeta \in D \cap V,\ X \in \mathbb{C}^n.$$

Set $t' = \sup\{t : s([0,t]) \subset V\}$ and $z' = s(t')$. Then

$$\begin{aligned} r' + \varepsilon > k_D(q,z) + \varepsilon &> \int_0^{t'} \kappa_D(s(t); s'(t))dt \\ &\geq p\int_0^{t'} \kappa_{D\cap U}(s(t); s'(t))dt \geq pk_{D\cap U}(q,z') \geq pc_{D\cap U}(q,z'). \end{aligned}$$

Taking a peak function for $D \cap U$ at a as a competitor in the definition of $c_{D\cap U}$, it follows that

$$\liminf_{q\to a\ \zeta\notin V} c_{D\cap U}(q,\zeta) = +\infty.$$

Therefore, we may find a neighborhood $W \subset V$ such that if $q \in W$, then $z' \in V$. Therefore $z' = z$ and the claim follows by letting $\varepsilon \to 0$.

(iii) In this case we do not know if a is a local peak point. However, Theorem 1 (iii) together with two small modifications in the previous proof imply the desired result.

1) The strong localization for the Kobayashi metric follows by the property

$$\liminf_{q\to a\ \zeta\notin\mathcal{U}} l_D(q,\zeta) = +\infty,$$

where $\mathcal{U}$ is any neighborhood of a (cf. [5, Proposition 7.2.9]).

It is easy to see that this property is a consequence of the following

Claim 2. If $(\varphi_j) \subset \mathcal{O}(\mathbb{D}, D)$ and $\varphi_j(0) \to a$, then $\varphi_j \rightrightarrows a$.

To prove Claim 2, assume the contrary. Since D is bounded, then, passing to a subsequence (if necessary), we may suppose that $\varphi_j \rightrightarrows \varphi \in \mathcal{O}(\mathbb{D}, \overline{D})$ and $\varphi \neq a$. Using again that D is bounded, we may find an $s \in (0,1)$ such that $\varphi_j(s\mathbb{D}) \subset U$ for any j. Note now that [10, Proposition 3] (see also [11, Proposition 1.5]) implies the tautness of bounded $\mathbb{C}$-convex domains. Then $D \cap U$ is a taut domain and hence

$\varphi(s\mathbb{D}) \in \partial D$. Since ∂D contains no affine discs through $a \in \partial D$, we get similarly to the proof of [12, Proposition 7] that $\varphi(s\mathbb{D}) = \{a\}$. Then the identity principle implies that $\varphi = a$. This contradiction completes the proof of Claim 2.

2) $\liminf\limits_{q\to a\ \zeta\notin V} k_{D\cap U}(q,\zeta) = +\infty$.[4]

This follows by Theorem 1 (iii) and the equality (2).

References

[1] M. Andersson, M. Passare, R. Sigurdsson, *Complex convexity and analytic functionals*, Birkhäuser, Basel-Boston-Berlin, 2004.

[2] G. Aladro, *The comparability of the Kobayashi approach region and the admissible approach region*, Illinois J. Math. 33 (1989), 42-63.

[3] G. P. Balakumar, P. Mahajan, K. Verma, *Bounds for invariant distances on pseudoconvex Levi corank nne domains and applications*, manuscripta math. (to appear); arXiv:1303.3439.

[4] M. Conrad, *Nicht isotrope Abschätzungen für lineal konvexe Gebiete endlichen Typs*, Dissertation, Universität Wuppertal, 2002.

[5] M. Jarnicki, P. Pflug, *Invariant distances and metrics in complex analysis*, de Gruyter, Berlin-New York (1993).

[6] P. Mahajan, K. Verma, *Some aspects of the Kobayashi and Carathéodory metrics on pseudoconvex domains*, J. Geom. Anal. 22 (2012), 491-560.

[7] N. Nikolov, *Localization of invariant metrics*, Arch. Math. 79 (2002) 67-73.

[8] N. Nikolov, P. Pflug, *Behavior of the Bergman kernel and metric near convex boundary points*, Proc. Amer. Math. Soc. 131 (2003), 2097-2102.

[9] N. Nikolov, P. Pflug, *Estimates for the Bergman kernel and metric of convex domains in* $\mathbb{C}^n$, Ann. Polon. Math. 81 (2003), 73-78.

[10] N. Nikolov, P. Pflug, W. Zwonek, *An example of a bounded* $\mathbb{C}$*-convex domain which is not biholomorphic to a convex domain*, Math. Scand. 102 (2008), 149-155.

[11] N. Nikolov, P. Pflug, P. J. Thomas, W. Zwonek, *On a local characterization of pseudoconvex domains*, Indiana Univ. Math. J. 58 (2009), 2661-2671.

[12] N. Nikolov, P. Pflug, W. Zwonek, *Estimates for invariant metrics on* $\mathbb{C}^n$*-convex domains*, Trans. Amer. Math. Soc. 363 (2011), 6245-6256.

[13] N. Nikolov, *Estimates of invariant metrics on "convex domains"*, Ann. Mat. Pura Appl., DOI 10.1007/s10231-013-0345-7.

[14] S. Sahin, *Poletsky–Stessin Hardy spaces on complex ellipsoids in* $\mathbb{C}^n$, arXiv: 1403.2506.

Institute of Mathematics and Informatics, Bulgarian Academy of Sciences, Acad. G. Bonchev 8, 1113 Sofia, Bulgaria

Faculty of Information Sciences, State University of Library Studies and Information Technologies, Shipchenski prohod 69A, 1574 Sofia, Bulgaria

E-mail address: nik@math.bas.bg

[4] This equality instead of the same for $c_{D\cap U}$ can be used to finish the proof of Proposition 3 (ii).

Institute of Mathematics, Faculty of Mathematics and Computer Science,, Jagiellonian University, Łojasiewicza 6, 30-348 Kraków, Poland
E-mail address: maria.trybula@im.uj.edu.pl